

## UCRL-JC-120470 Abs

### **The Effect of Biomass Burning on Tropospheric Ozone Production: Three-Dimensional, Global Modeling Results\***

C.S. Atherton, J.E. Penner, J.J. Walton (Global Climate Research Division, Lawrence Livermore National Laboratory, P.O. Box 808, Livermore, CA 94551; e-mail: cyndi@tropos.llnl.gov)

Tropospheric ozone (O<sub>3</sub>) is formed when CO, CH<sub>4</sub>, and non-methane hydrocarbons (NMHCs) react in the presence of sunlight and NO<sub>x</sub>. In the tropics and southern hemisphere, biomass burning is a significant source of many O<sub>3</sub> precursors. Thus, biomass burning may strongly affect levels of O<sub>3</sub> and other species in many regions of the world.

We have developed a three-dimensional, global chemistry-transport-deposition model with 76 prognostic chemical species (including O<sub>3</sub>, OH, PAN, NO<sub>x</sub>, HNO<sub>3</sub>, isoprene, and a full suite of non-methane hydrocarbons). Here we present results of both a baseline simulation, and a simulation in which biomass burning emissions are significantly reduced. The results show the extent and magnitude to which biomass burning emissions affect tropospheric O<sub>3</sub> production on a global scale.

\*This work was performed under the auspices of the U.S. Department of Energy by the Lawrence Livermore National Laboratory under Contract No. W-7405-Eng-48.

## Submittal Information

1. 1995 IUGG Meeting
2. 007724433
3. (a) Cynthia Atherton  
LLNL  
P.O. Box 808, L-262  
Livermore, CA 94551  
(b) Tel:510-422-1825  
(c) FAX:510-422-5844  
(d) D-Mail: cyndi@tropos.llnl.gov
4. IAMAS
5. M1 - Tropospheric Ozone
6. Oral
7. None
8. No